

HARMFUL
KEEP OUT OF REACH OF CHILDREN



Trace-It *Copper*TM

**Liquid Copper Chelate for the correction
of Copper deficiencies**

**Contains: 7.0% w/v (70g/L) Copper (Cu) – expressed
as elemental content**



MANUFACTURED & DISTRIBUTED BY:

AgriNova NZ Ltd
15 Sunlight Grove, PO Box 50-080
Porirua, New Zealand
Freephone NZ 0800 447 624
Freephone AUS 1800 777 068
www.grochem.co.nz



9142102611716891

Approved Pursuant to the HSNO Act 1996, under the Fertilisers – Subsidiary Hazard group standard HSR2571.
See www.epa.govt.nz for approval conditions.

Net Contents: 20 Litres

PLEASE READ LABEL AND SAFETY DATA SHEET COMPLETELY BEFORE USE

PRECAUTIONS:

Keep out of reach of children. Store in original container, away from foodstuffs. Wear suitable impervious gloves and eye protection and avoid contact with any exposed skin. Wash hands with warm soapy water after use and before handling food. In case of eye contact, flush with plenty of water for 15 minutes, and seek medical attention if irritation persists.

STORAGE

Keep in cool, dry chemical store, out of reach of children and animals. Protect from frost and other climatic extremes (including temperatures exceeding 40°C). Poor water quality may reduce the effectiveness of the product, the solubility of the product and/ or increase the potential for phytotoxic effects on plants.

CONDITIONS OF SALE:

As no control can be exercised over the manner in which or the conditions under which this product is used, all conditions and warranties expressed or implied, other than those required by statute are excluded. No responsibility will be accepted for any damage or injury whatsoever arising from the storage, handling, application or use of this product.

Batch:

DoM:

DIRECTIONS FOR USE

ALWAYS FOLLOW LABEL INSTRUCTIONS.

- Mix Trace-it Copper™ in its container thoroughly before use. Fill the spray tank until three quarters full with water and then, whilst agitating the spray tank, add the Trace-it Copper™ via the top filter. Continue agitation until application complete. Always apply diluted or mixed chemicals immediately, do not store.
- Tank mixing or co-application. Read all labels carefully and adhere strictly to instructions on mixing. Apply to small test area only when using unfamiliar products or new crops. Many variables are outside the control of Agri-Nova and can influence results of co-applied products and therefore co-application is entirely at the risk of the end user.

- Trace-it Copper™ should be applied in low light conditions in the morning or evening or on cloudy days. Do not apply during periods of high sunlight, heat, during rain or prior to expected rain.
- Where a deficiency is severe, additional applications may be required.

Trace-it Copper™ is specially designed as a high availability copper source to rapidly correct nutrient deficiencies in a wide range of crops. Copper is an essential micronutrient particularly active in many of the metabolic pathways of cells. Photosynthesis, respiration, cell wall permeability, certain proteins, nitrogen and carbohydrate processes are all affected by the availability of copper.

Crop	Rate per ha	Min water rate per ha	Recommendations
Kiwifruit	1-1.5 L	500L	As required, ensure good coverage.
Grapes	0.4-0.8 L	500L	As required ensure good coverage.
Fruit Crops	0.4-0.8 L	500L	As required ensure good coverage.
Field Crops	0.4-0.8 L	100L	As required, ensure good coverage.
Glass house	40-80ml per 100L water	50-100L per 1000m ²	As required, avoid application directly to flowers or material sensitive to marking.
Maize/Corn	400ml	200L	4-6 leaf stage.