

**HARMFUL: KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING**



GIB-47™

PLANT GROWTH REGULATOR

GIB-47 IMPROVES FRUIT QUALITY BY REDUCING THE INCIDENCE OF RUSSET AND IMPROVING SKIN FINISH.

grochem

Contains: 10g/litre Gibberellin A4/7 in the form of a soluble concentrate.

Batch:

DoM:

NET CONTENTS

5L

GGIB47/0617

REGISTERED TO & DISTRIBUTED BY: AgriNova New Zealand Ltd,
15 Sunlight Grove, PO Box 50-080, Porirua,
New Zealand. NZ
0800 447 624, AUS 1800 777 068
www.grochem.com



9 14 2 1 0 2 6 1 1 7 1 4 8 1

**HARMFUL: KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING**



GIB-47™

PLANT GROWTH REGULATOR

GIB-47 IMPROVES FRUIT QUALITY BY REDUCING THE INCIDENCE OF RUSSET AND IMPROVING SKIN FINISH.

grochem

Contains: 10g/litre Gibberellin A4/7 in the form of a soluble concentrate.

Batch:

DoM:

NET CONTENTS

5L

GGIB47/0617

REGISTERED TO & DISTRIBUTED BY: AgriNova New Zealand Ltd,
15 Sunlight Grove, PO Box 50-080, Porirua,
New Zealand. NZ
0800 447 624, AUS 1800 777 068
www.grochem.com



9 14 2 1 0 2 6 1 1 7 1 4 8 1

THIS PRODUCT MUST NOT BE USED FOR ANY PURPOSE OR ANY MANNER CONTRARY TO THIS LABEL

FIRST AID: IF SWALLOWED: If large quantity is swallowed, give lukewarm water (½ litre) if victim is completely conscious/alert. Obtain medical attention.
IF IN EYE: Thoroughly flush eyes with large amounts of clean water for at least 15 minutes. If irritation persists, seek medical attention.
IF ON SKIN: Wash with soap and water.
IF INHALED: Remove victim to fresh air. Obtain medical attention if symptoms persist.

24-hour Emergency Telephone Number – For advice contact the National Poisons Centre – 0800 POISON (0800 764 766) – or a doctor, immediately. Have product container or label at hand.

For specialist advice in an emergency only, call CHEMCALL (24hrs) 0800 243 6225. In a transport emergency dial 111.

SAFETY DATA SHEET: A Safety Data Sheet (SDS) should be available at all work places where Gib-47 is stored or used.

Additional copies are available from Grochem on request or www.grochem.com

STORAGE: Store in a cool location away from foodstuffs. Avoid all sources of ignition. When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

DISPOSAL: Product disposal: Use as directed or dispose of at an approved landfill.

Container disposal: Triple rinse container and crush before disposal at an approved landfill.

EQUIPMENT: Apply using accurately calibrated and maintained equipment in accordance with The New Zealand Standard for the Management of Agrochemicals (NZS8409).

SPILLAGE: Absorb spillage with sawdust or other absorbent material. Rinse area with water and absorb rinsate.

DIRECTIONS FOR USE: Gib-47 is used to reduce russet in russet prone apple varieties and reduce splitting in Nashi and cherries. Gib-47 is believed to act by increasing elasticity and production of the outer epidermal cells. Russetting in apples is associated with skin damage due to prolonged wetness, low temperatures or imbalances in skin growth during the early season.

WARNINGS: Exceeding recommended rates is not recommended as higher than label rates may reduce the flowering in the following season.

COMPATIBILITY: Apply alone. May be applied with non-ionic surfactants.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residues Limits of Agricultural Compounds) Food Standards.

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

APPLICATION RATES:

CROP	RUSSET PREVENTION	RATE
Coxs Orange		Apply 50ml
Fuji, Royal Gala, Pacific Rose and other varieties		Apply 50–100ml
Miscellaneous Uses: When applied to a wide range of annual and perennial crops and ornamentals this product is known to be effective in many applications including: stem elongation, breaking of seed dormancy, earlier flowering and setting of fruit under adverse weather conditions. Examples: Nashi – (Nijisseiki) and Cherries.		Apply 100ml–1L

WATER RATE: Apply 1000L of water/ha.

On apples make the first application at late flowering and in advance of poor weather conditions. Make two or three further applications at 7–10 day intervals.

Use the higher rate and greater numbers of applications where the risk of russetting is higher.
Apply with a calibrated sprayer that ensures uniform and complete coverage.

High relative humidity and slow drying conditions favour maximum absorption of Gib-47.

Rainfall within 6 hours may reduce the activity.

CONDITIONS OF SALE: As no control can be exercised over the manner in which or the conditions under which this product is used, all conditions and warranties expressed or implied, other than those required by statute are excluded. No responsibility will be accepted for any damage or injury whatsoever arising from the storage, handling, application or use of this product.

Registered pursuant to the ACVM Act 1997, No. P7228. See www.foodsafety.govt.nz for registration conditions.

THIS PRODUCT MUST NOT BE USED FOR ANY PURPOSE OR ANY MANNER CONTRARY TO THIS LABEL

FIRST AID: IF SWALLOWED: If large quantity is swallowed, give lukewarm water (½ litre) if victim is completely conscious/alert. Obtain medical attention.
IF IN EYE: Thoroughly flush eyes with large amounts of clean water for at least 15 minutes. If irritation persists, seek medical attention.
IF ON SKIN: Wash with soap and water.
IF INHALED: Remove victim to fresh air. Obtain medical attention if symptoms persist.

24-hour Emergency Telephone Number – For advice contact the National Poisons Centre – 0800 POISON (0800 764 766) – or a doctor, immediately. Have product container or label at hand.

For specialist advice in an emergency only, call CHEMCALL (24hrs) 0800 243 6225. In a transport emergency dial 111.

SAFETY DATA SHEET: A Safety Data Sheet (SDS) should be available at all work places where Gib-47 is stored or used.

Additional copies are available from Grochem on request or www.grochem.com

STORAGE: Store in a cool location away from foodstuffs. Avoid all sources of ignition. When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

DISPOSAL: Product disposal: Use as directed or dispose of at an approved landfill.

Container disposal: Triple rinse container and crush before disposal at an approved landfill.

EQUIPMENT: Apply using accurately calibrated and maintained equipment in accordance with The New Zealand Standard for the Management of Agrochemicals (NZS8409).

SPILLAGE: Absorb spillage with sawdust or other absorbent material. Rinse area with water and absorb rinsate.

DIRECTIONS FOR USE: Gib-47 is used to reduce russet in russet prone apple varieties and reduce splitting in Nashi and cherries. Gib-47 is believed to act by increasing elasticity and production of the outer epidermal cells. Russetting in apples is associated with skin damage due to prolonged wetness, low temperatures or imbalances in skin growth during the early season.

WARNINGS: Exceeding recommended rates is not recommended as higher than label rates may reduce the flowering in the following season.

COMPATIBILITY: Apply alone. May be applied with non-ionic surfactants.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residues Limits of Agricultural Compounds) Food Standards.

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

APPLICATION RATES:

CROP	RUSSET PREVENTION	RATE
Coxs Orange		Apply 50ml
Fuji, Royal Gala, Pacific Rose and other varieties		Apply 50–100ml
Miscellaneous Uses: When applied to a wide range of annual and perennial crops and ornamentals this product is known to be effective in many applications including: stem elongation, breaking of seed dormancy, earlier flowering and setting of fruit under adverse weather conditions. Examples: Nashi – (Nijisseiki) and Cherries.		Apply 100ml–1L

WATER RATE: Apply 1000L of water/ha.

On apples make the first application at late flowering and in advance of poor weather conditions. Make two or three further applications at 7–10 day intervals.

Use the higher rate and greater numbers of applications where the risk of russetting is higher.
Apply with a calibrated sprayer that ensures uniform and complete coverage.

High relative humidity and slow drying conditions favour maximum absorption of Gib-47.

Rainfall within 6 hours may reduce the activity.

CONDITIONS OF SALE: As no control can be exercised over the manner in which or the conditions under which this product is used, all conditions and warranties expressed or implied, other than those required by statute are excluded. No responsibility will be accepted for any damage or injury whatsoever arising from the storage, handling, application or use of this product.

Registered pursuant to the ACVM Act 1997, No. P7228. See www.foodsafety.govt.nz for registration conditions.