



SOLETO™

Pre-emergent herbicide for potatoes. European technology for New Zealand potato growers.

- · Active Metobromuron 500g/LSC
- · Activity Photosynthesis inhibitor (II)
- Resistance group C2 Herbicide
- Mode of action Contact and residual activity.
- Weed target Broadleaf weeds (Dicots) and some grasses.
- Current NZ label Annual broadleaf weeds, Redroot and Fathen

PRODUCT BENEFITS

- Broad weed spectrum.
- Very high crop safety, no variety restrictions.
 Safe on Metribuzin sensitive varieties such as Innovator and VR808.
- Good persistence half life of 22 days in soil and only moderately soluble, ensuring effective weed control until after row closure.
- Effective under all conditions with consistent result. Including drought conditions.
- Tank mix flexibility talk to your local agronomist for recommendations.
- Tried and trusted in UK and Europe for more than 10 years.
- Assessed for efficacy and crop safety in multiple trials in Canterbury and Pukekohe over three years prior to registration.

APPLICATION TIMING

Pre-emergence.

PRODUCT RATE

3 L per hectare when applied alone. Discuss tank mix combinations with your agronomist.

Registered pursuant to the ACVM Act 1997, No. P9699. See www.foodsafety.govt.nz for registration conditions. Approved Pursuant to the HSNO Act 1996, Number HSR 101457. See www.epa.govt.nz for approval conditions.





DISCLAIMER: The information contained in this publication is of a general nature and should not be relied upon as a substitute for professional advice in specific cases. As no control can be exercised over the manner in which or the conditions under which this product is used, no responsibility will be accepted for any damage or injury whatsoever arising from the storage, handling, application or use of this product.

